

HEXAMITE® 1AQ SELECT

Miticide for use on turf, ornamentals and non-bearing fruit trees and vines

ACTIVE INGREDIENT

Hexythiazox – Group 10A

% ACTIVE INGREDIENT

11.8%

FORMULATION

Aqueous suspension

CHEMICAL FAMILY

Thiazolidinone

BRAND ALTERNATIVE

Hexygon® IQ

MODE OF ACTION

Controls mite growth through activity on eggs and early stages of development.

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

12 x 1 qt

LABELLED USES

Turf

Ornamentals

In and around residential, institutional, public, commercial, agricultural and industrial buildings



Hexamite® 1AQ Select controls mites through activity on eggs and immature stages through either direct contact with the spray or from contact with treated plant surfaces. Its ovicidal activity controls newly deposited mite eggs and eggs that are laid after application. Also highly effective in controlling immature motile stages of target mite species that have been sprayed or have moved onto treated surfaces. Adult mites are not directly affected. The degree and duration of control is dependent on the rate used, growth stage of the mite, species of mite, and climatic conditions under which the material is applied.

PRODUCT BENEFITS

- Long lasting miticide with residual control for up to two months
- Unique mode of action provides effective activity on eggs and early stages of development
- For use on turf, ornamentals, fruit trees and vines



Prime Source, a Division of Albaugh, LLC

1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2022 Albaugh, LLC. Prime Source, the water drop logo and Hexamite are registered trademarks of Albaugh, LLC. Hexygon is a registered trademark of Gowan Company. Always read and follow label Precautions and Directions For Use.

HEXAMITE® 1AQ SELECT

Miticide for use on turf, ornamentals and non-bearing fruit trees and vines

Labeled mite pests for turf, ornamentals and non-bearing fruit trees and vines

Arborvitae spider mite	European red mite	Spruce spider mite	Two-spotted spider mite
Banks grass mite	Honeylocust spider mite	Strawberry spider mite	Willamette mite
Brown wheat mite	Pacific spider mite	Tetranychid mites	
Citrus red mite	Pecan leaf scorch mite	(such as European red mite	
Clover mite	Southern red mite	and two-spotted spider mite)	

Application directions for ornamentals and turf

CROP	PEST	RATE	USE DIRECTIONS
ORNAMENTAL PLANTS AND NON-BEARING FRUIT TREES AND VINES Growing in nurseries, greenhouses, shadehouses and Christmas tree plantations	Arborvitae spider mite European red mite Honeylocust spider mite Pacific spider mite Southern red mite Spruce spider mite Strawberry spider mite Two-spotted spider mite Willamette mite Banks Grass mite Clover mite Brown wheat mite	12 – 24 ozs/acre OR 4 – 8 ozs/100 gal OR 0.28 – 0.55 ozs/ 1000 sq ft	When using high-volume dilute sprays, do not exceed the maximum per acre rate or apply less than the minimum labeled rate. For concentrate equipment, use the recommended rate per acre. Apply to lawns and turf at the first signs of egg deposition.
ESTABLISHED ORNAMENTAL LANDSCAPE PLANTINGS Including those in: Interiorscapes, residences, public areas, commercial areas, institutional areas, industrial areas, rights of way and other easements, and recreational sites such as campgrounds, golfcourses, parks and athletic fields			
ORNAMENTAL LAWNS AND TURF (including residential) Except California			
NON-RESIDENTIAL TURFGRASS (including turf grown for sod, turf grown to produce seed, or turf for research purposes) Except California			

See label for full list of pests controlled and application rates.

Prime Source, a Division of Albaugh, LLC • 1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2022 Albaugh, LLC. Prime Source, the water drop logo and Hexamite are registered trademarks of Albaugh, LLC. Hexygon is a registered trademark of Gowan Company. Always read and follow label Precautions and Directions For Use.